



Ring Ouzel. Adult. Male (05-X).

### **IDENTIFICATION**

24-27 cm. Adult male black in breeding plumage; pectoral band pure white; yellow bill with dark tip; in non breeding plumage in winter with greyish edges on feathers and dark bill with yellowish tip. Female with dark grey plumage; pectoral band dark brown. Some 1st year females without pale pectoral band.





Ring Ouzel. Pattern of wing, tail and upperparts.



### SIMILAR SPECIES

The white pectoral band makes this species unmistakable. Females without white band recall a Spotless Starling with has shorter tail and thinner bill; Female Blue Rock Thrush has similar size, but different pattern on flanks and underparts; Blackbird has wing feathers without pale edges.









### **SEXING**

In adult birds, breeding male with a white or light brownish pectoral band; head and neck black or brownish-black; in autumn feathers on head and neck with brownish edges; body feathers with a narrow pale edge (about 1 mm). Breeding female with obscure brownish-white pectoral band; head and neck dak brownish; in autumn feathers on head and neck with paler edges; body feathers with a wide brownish edge (about 2 mm) (CAUTION: some adult females are very similar to males). In 1st year autumn/2nd year spring birds, male with an evident pectoral band, paler than underparts; female with poor marked pectoral band, with the same colour than underparts. Juveniles cannot be sexed.





Ring Ouzel. Adult. Sexing. Pattern of band on breast: left male; right female.





Ring Ouzel. 1st year. Sexing. Pattern of band on breast: left male; right female.

### **AGEING**

3 types of age can be recognized:

**Juvenile** has body feathers with pale tips; pale chin and throat; pale mottled on tips of lesser and median coverts; greater coverts with large whitish edge.

1st year autumn/2ne year spring with moult limit between the inner changed greater coverts (narrow pale edge) and the outermost unmoulted; tail feathers with pointed tips; some females with unmarked pectoral band.

**Adult** with no moult limits on wing; all the greater coverts with narrow edge; tail feathers with rounded tips.





Ring Ouzel. Ageing. Pattern of breast: left adult female; right juvenile (photo: Alex Fonollosa).





Ring Ouzel. Ageing. Pattern of wing coverts: top adult female; bottom juvenile (photo: Alex Fonollosa).







Ring Ouzel. Ageing. Pattern of tips of tail feathers: left adult; right 1st year.



Ring Ouzel. Male. Ageing. Moult limit on wing coverts: top adult; bottom 1st year.






Ring Ouzel. Female. Ageing. Moult limit on wing coverts: top adult; bottom 1st year.

### **MOULT**

Complete **postbreeding** moult, usually finished in September before migration. Partial **postjuvenile** moult including body feathers, lesser and median coverts and a variable number of greater coverts; some birds can moult tertials; usually finished in October.



Ring Ouzel. Extent of postjuveni-le moult.

### **PHENOLOGY**







Ring Ouzel. Adult. Female (10-I).






Ring Ouzel.
Adult. Head
pattern: top
male (05-X);
bottom female
(10-I).



Ring Ouzel. 1st year. Male (01-XI).







Ring Ouzel. 1st year. Female (07-X).








Ring Ouzel. Juvenile (14-VII) (photo: Alex Fonollosa).







Ring Ouzel. Adult. Crown pattern: left male (05-X); right female (10-I).







Ring Ouzel. Crown pattern: top left male (01-XI); top right female (07-X); left juvenile (14-VII) (photo: Alex Fonollosa).






Ring Ouzel. Adult. Neck pattern: left male (05-X); right female (10-I).











Ring Ouzel. Adult. Breast pattern: left male (05-X); right female (10-I).







Ring Ouzel. Breast pattern: top left male (01-XI); top right female (07-X); left juvenile (14-VII) (photo: Alex Fonollosa).

http://blascozumeta.com

http://www.aranzadi.eus







Ring Ouzel. Adult. Upperparts pattern: left male (05-X); right female (10-I).







Ring Ouzel. Upperparts pattern: top left male (01-XI); top right female (07-X); left juvenile (14-VII) (photo: Alex Fonollosa).





Ring Ouzel. Adult. Tail pattern: left male (05-X); right female (10-I).







Ring Ouzel. Tail pattern: top left male (01-XI); top right female (07-X); left juvenile (14-VII) (photo: Alex Fonollosa).



Ring Ouzel. Adult. Male: pattern of wing (05-X).

http://blascozumeta.com http://www.aranzadi.eus 1





Ring Ouzel. Adult. Female: pattern of wing (10-I).



Ring Ouzel. 1st year. Male: pattern of wing (01-XI).



Ring Ouzel. 1st year. Female: pattern of wing (07-X).



Ring Ouzel. Juvenile: wing pttern (14-VII) (photo: Alex Fonollosa).